

First-order scaling near a second-order phase transition: Tricritical polymer collapse

A. L. OWCZAREK^{1(*)} and T. PRELLBERG^{2(**)}

¹ *Department of Mathematics and Statistics, The University of Melbourne
3010, Melbourne, Australia*

² *Department of Physics, Syracuse University
Syracuse, NY 13244, USA*

(received 8 February 2000; accepted in final form 28 July 2000)

PACS. 05.50.+q – Lattice theory and statistics.

PACS. 05.70.Fh – Phase transitions: general studies.

PACS. 61.41.+e – Polymers, elastomers, and plastics.

Abstract. – The coil-globule transition of an isolated polymer has been well established to be a second-order phase transition described by a standard tricritical $O(0)$ field theory. We provide compelling evidence from Monte Carlo simulations in four dimensions, where mean-field theory should apply, that the approach to this (tri)critical point is dominated by the build-up of first-order-like singularities masking the second-order nature of the coil-globule transition: the distribution of the internal energy having two clear peaks that become more distinct and sharp as the tricritical point is approached. However, the distance between the peaks slowly decays to zero. The evidence shows that the position of this (pseudo) first-order transition is shifted by an amount from the tricritical point that is asymptotically much larger than the width of the transition region.

An isolated polymer in solution is usually argued to be in one of three states depending on the strength of the inter-monomer interactions which are mediated by the solvent molecules and can be controlled via the temperature T . Let us define the radius of gyration exponent ν as

$$R_N \sim N^\nu \quad \text{as} \quad N \rightarrow \infty, \quad (1)$$

where R_N is the radius of gyration. Note that confluent logarithmic factors may also appear in this form. At high temperatures and in so-called “good solvents” a polymer chain is expected to be in a swollen phase (swollen coil) relative to a reference Gaussian state so that the average size of the polymer, as measured by the radius of gyration, scales with chain length faster than it would if it were behaving as a random walk. In dimensions 2 and 3 it is expected that the swollen value of $\nu = \nu_s > 1/2$. At low temperatures or in poor solvents the polymer is expected to be in a collapsed globular form with a macroscopic density inside the polymer: this implying an average size that scales slower [1] than a random walk, in particular

(*) E-mail: aleks@ms.unimelb.edu.au

(**) E-mail: tprell@physics.syr.edu

$\nu = \nu_g = 1/d < 1/2$. Between these two states there is expected to be a second-order phase transition (sharp in the infinite chain length limit). The standard description of the collapse transition is that of a tricritical point related to the $n \rightarrow 0$ limit of the ϕ^4 – ϕ^6 $O(n)$ field theory [1–3]. One then might expect that above the upper critical dimension ($d_u = 3$) some type of self-consistent mean-field theory based upon a suitable tricritical Landau-Ginzberg Hamiltonian [1, 4] would give a full description of the transition, and hence conclude that in all dimensions $d > 3$ there is a collapse transition from a swollen state to the globular state with classical tricritical behaviour. The classical theory predicts that at the transition point the polymer actually behaves as if it were a random walk ($\nu = \nu_\theta = 1/2$), and this point $T = T_\theta$ has been known as the θ -point.

Around the transition ($T \rightarrow T_\theta$ and $N \rightarrow \infty$) a tricritical crossover scaling form [5] describing the rounding of the transition is generally predicted to be

$$R_N(T) \sim N^{\nu_\theta} \mathcal{F}((T_\theta - T)N^\phi), \quad (2)$$

where the scaling function $\mathcal{F}(z)$ is a strictly positive analytic function of z (note that $\mathcal{F}(0) > 0$). The exponent ϕ is known as the crossover exponent. Moreover, the asymptotic behaviour of $\mathcal{F}(z)$ for $z \rightarrow \pm\infty$ is expected to “match” the asymptotic behaviour of $R_N(T)$. For example, consider fixing T such that $T < T_\theta$ so that $R_N(T)$ obeys (1) with $\nu = 1/d$. It follows that the scaling function $\mathcal{F}(z)$ must satisfy

$$\mathcal{F}(z) \sim z^{(1/d - \nu_\theta)/\phi} \quad \text{as } z \rightarrow \infty. \quad (3)$$

Generically, it is expected that forms such as (2) provide all the information needed about the scaling around the tricritical point. In particular, if one measures the “shift”, $T_{c,N} - T_\theta$, of the transition at finite lengths by, say, finding the position of a peak in the specific heat, $T_{c,N}$, or other appropriate derivative of the free energy, then the associated exponent ψ_p , defined by

$$T_{c,N} - T_\theta \sim N^{-\psi_p} \quad \text{as } N \rightarrow \infty, \quad (4)$$

should obey the relation $\psi_p = \phi$.

The application of the mean-field theory of a tricritical point to polymer collapse predicts a weak transition in the thermodynamic limit, with a jump in the specific heat, that is, $\alpha = 0$. In three dimensions the application of various self-consistent mean-field-like approaches predicts that the second-order transition is rounded and shifted on the same scale of $N^{-1/2}$, that is, $\phi = \psi_p = 1/2$, though strictly the power laws involved are modified via renormalisation group arguments by confluent logarithms. In four and higher dimensions no confluent logarithms should be present and one may expect pure mean-field behaviour with a crossover exponent of $1/2$. However the following simple scaling argument may be invoked. Consider the tricritical scaling form (2) with $\nu_\theta = 1/2$, and especially consider the behaviour of (2) for $T < T_\theta$ and T fixed. As stated above, to match the fixed- T behaviour of $R_N(T)$ the scaling function $\mathcal{F}(z)$ must obey (3). Now define the internal density of a polymer chain ρ as $\rho = N/R^d$ since R^d is the approximate volume the polymer assumes. In the thermodynamic limit $\rho = 0$ for $T > T_\theta$ and it is expected that $\rho \sim a(T_\theta - T)^\beta$ as $T \rightarrow T_\theta^-$. In mean-field theory the exponent $\beta = 1$. Using this and $\nu_\theta = 1/2$ one can deduce that $\phi = (d/2 - 1)$. Of course this is in contradiction to the assumption above that $\phi = 1/2$ and leads one to suspect that there may be something unexpected happening in dimensions $d > 3$.

To consider such issues we have simulated interacting self-avoiding walks, the canonical lattice model of polymer collapse, on the four-dimensional hyper-cubic lattice using the PERM algorithm [6] over a wide range of temperatures with surprising results. Because this algorithm




Fig. 1 – Internal energy density distributions for lengths $N = 2516811684$ and $N = 2516811684$ calculated by the method of [1] and [2].




Fig. 2 – Scaling of the shift of the transition: we show the scaling combination $N^{1/3}(\omega_{c,N} - \omega_\theta)$ vs. $N^{-2/3}$, where $\omega_{c,N}$ and ω_θ are the Boltzmann weights associated with the monomer-monomer interaction at $T_{c,N}$ and T_θ .

dependence on the distance to the θ -point and so an exponent $\alpha = 0$, implying that a second-order phase transition occurs in the thermodynamic limit. From the theory one finds a rounded transition for finite N occurring at temperature $T_{c,N} < T_\theta$ with *shift* exponent

$$\psi_p = 1/(d-1), \quad (5)$$

so that the collapse occurs at finite length at a temperature that scales towards the θ -point quite slowly and is below the θ -point. This then concurs with our finding that $\psi_p \approx 1/3$ in $d = 4$. The width of this transition ΔT at finite N can be found and has *crossover* exponent

$$\phi = (d-2)/(d-1). \quad (6)$$

All the exponents are derived from the assumptions of mean-field thermodynamic behaviour and using $\nu_g = 1/d$. Hence all the exponents quoted here are related to each other (only one independent exponent). Our simulations show just such a narrow crossover region with a crossover exponent that is certainly larger than the shift exponent (the data is compatible with exponent $2/3$).

By considering the difference of the density ρ_s of the swollen state to the globular state ρ_g relative to the density of the swollen state at $T_{c,N}$, Khokhlov [8] concluded from its divergence that “the coil-globule transition is first-order”, though we now interpret this to mean that the finite-size corrections to the thermodynamic second-order transition are first-order-like. We point out that the terminology of Khokhlov was presumably that explained in sect. I.C.2 of [11] but may be misleading to the modern reader. However, both $\rho_g(T_{c,N})$ and $\rho_s(T_{c,N})$ tend to zero as $N \rightarrow \infty$ and it is simply that $\rho_g(T_{c,N})$ tends to zero asymptotically slower than $\rho_s(T_{c,N})$ that makes the relative difference diverge. The analysis can be used to deduce the scaling of R_N at $T_{c,N}$ with an exponent $\nu_c = 1/(d-1)$. Note that $\nu_\theta > \nu_c > \nu_g$, so that this scaling is in between the scaling fixed at the θ -point and at any temperature fixed in the collapsed phase. Following the work of Lifshitz, Grosberg and Khokhlov [11] one can also calculate the change in the internal energy over the crossover width of the transition ΔT as the latent heat ΔU from the free energy expression. The latent heat decays as N increases with exponent $1/(1-d)$. The corresponding height of the peak in the specific heat diverges with

exponent $(d-3)/(d-1)$. From our data we were unfortunately unable to extract reasonable estimates for ν_c or the exponents of the latent and specific heats.

To interpret our results, we can take the understanding of this mean-field theory further. Let us consider the distribution of internal energy, which we measured in our simulations, as a function of temperature and length. For any temperature above the θ -point and well below $T_{c,N}$ one expects the distribution of internal energy to look like a single-peaked distribution centred close to the thermodynamic limit value: a Gaussian distribution is expected around the peak with variance $O(N^{-1/2})$. In fact this picture should be valid for all temperatures outside a range of order $O(N^{-(d-2)/(d-1)})$ centred on $T_{c,N}$. When we enter this region we will expect to see a double-peaked distribution as in a first-order transition region. For any temperature in this transition region there should be two peaks in the internal energy distribution separated at the order of $O(N^{-1/(d-1)})$ (the value of the gap being the latent heat). Each of these peaks should be of Gaussian type with individual variances again of the order $O(N^{-1/2})$. Hence as N increases the peaks will become more and more distinct and relatively sharper but the peak positions will be getting closer together. Hence we refer to this scenario as a *pseudo*-first-order transition or, more correctly, as first-order-like finite-size corrections to a second-order phase transition. If there were a real first-order transition then the distance between the peaks should converge to a non-zero constant. A comprehensive interpretation of our computer simulations in $d=4$ [7] is most consistent with just such a scenario and leads us to conjecture that this theoretical picture is indeed correct for the coil-globule transition for $d \geq 4$. Preliminary simulational results for $d=5$, and for another model (interacting lattice trails) in $d=4$, demonstrate similar behaviour.

Let us return to the question of crossover scaling forms and our finding of two shift exponents from our computer simulations. While we cannot ascertain either with great accuracy, let us assume that we have a region around the θ -point that is approximated well by a form like (2) with crossover exponent $\phi_\theta = (d/2 - 1)$. Now, despite the fact that this does not describe the collapse transition region, we notice that substituting $t \sim N^{1/(d-1)}$, and using the asymptotics derived from our matching argument above, leads to $R \sim N^{1/(d-1)}$, which is precisely the correct scaling for the real transition region! Hence we conjecture a phenomenological *product* scaling form: (for $T < T_\theta$)

$$R_N \sim N^{1/2} \mathcal{F}((T_\theta - T)N^{\phi_\theta}) \mathcal{G}((T_{c,N} - T)N^\phi), \quad (7)$$

with $T_{c,N} \sim T_\theta - aN^{-\psi_p}$ and where $\mathcal{G}(y) \sim 1$ for $y \rightarrow \pm\infty$. This form will then correctly describe both the region around the θ -point and the rounded transition around $T_{c,N}$ and will match with the behaviour of the collapsed phase for fixed $T < T_\theta$. Such a form is not dependent on the finding of pseudo-first-order behaviour and may be useful for analysing data whenever two shift exponents are found.

In conclusion, Monte Carlo simulations of lattice polymers in four dimensions show that for finite length the rounded coil-globule transition appears to be first order but we argue that the tricritical predictions may well reappear in the infinite length limit as our results fit best the predictions of Lifshitz-Grosberg-Khokhlov (LGK) theory applied to high dimensions. We suggest that the crossover scaling forms are more complicated than at low dimensions and suggest a generalisation that may be heuristically useful.

* * *

Financial support from the Australian Research Council is gratefully acknowledged by ALO. This work was partially supported by EPSRC Grant No. GR/K79307. We thank A. J. GUTTMANN and the referees for many useful comments on the manuscript.

REFERENCES

- [1] DE GENNES P.-G., *J. Phys. (Paris) Lett.*, **36** (1975) L55.
- [2] DE GENNES P.-G., *J. Phys. (Paris) Lett.*, **39** (1978) L299.
- [3] DUPLANTIER B., *J. Phys. (Paris)*, **43** (1982) 991.
- [4] LAWRIE I. D. and SARBACH S., in *Phase Transitions and Critical Phenomena*, edited by DOMB C. and LEBOWITZ J. L., Vol. 9 (Academic, London) 1984.
- [5] DE GENNES P.-G., *Scaling Concepts in Polymer Physics* (Cornell University Press, Ithaca) 1979.
- [6] GRASSBERGER P., *Phys. Rev. E*, **56** (1997) 3682.
- [7] PRELLBERG T. and OWCZAREK A. L., unpublished.
- [8] KHOKHLOV A. R., *Physica A*, **105** (1981) 357.
- [9] LIFSHITZ I. M., *Zh. Eksp. Teor. Fiz.*, **55** (1968) 2408.
- [10] LIFSHITZ I. M., GROSBERG A. Y. and KHOKHLOV A. R., *Zh. Eksp. Teor. Fiz.*, **71** (1976) 1634.
- [11] LIFSHITZ I. M., GROSBERG A. Y. and KHOKHLOV A. R., *Rev. Mod. Phys.*, **50** (1978) 683.
- [12] OWCZAREK A. L., PRELLBERG T. and BRAK R., *Phys. Rev. Lett.*, **70** (1993) 951.
- [13] OWCZAREK A. L., *J. Phys. A*, **26** (1993) L647.
- [14] GRASSBERGER P. and HEGGER R., *J. Chem. Phys.*, **102** (1995) 6881.
- [15] NIDRAS P. P., *J. Phys. A*, **29** (1996) 7929.